

‘Host of the month’ is a series of information sheets and blogs that highlight a tree host and their associated priority pests and diseases that are best seen and recorded in that month. For September we’re looking at Planes (*Platanus* species), Canker stain of plane, and Plane lacebug.

The tallest broadleaf tree in the UK is a London plane (*Platanus x hispanica*) which grows in the grounds of Bryanston School in Dorset and was measured in 2015 at 49.7m. The origins of London plane (*Platanus x hispanica*), a hybrid between Oriental (*P. orientalis*) and Western plane (*P. occidentalis*), are uncertain but most likely happened in the 1600’s in Spain. It was first planted in England around 1680 at Ely in Cambridgeshire, and at Barnes in Surrey and both trees are still there today, the latter one going by the name of Barnie. By the 1920’s London plane represented around 60% of the trees in London though that has now dropped to around 4%.

London planes are still common in cities and parks across Europe, Asia, South Africa and North America, in part because they are adept at trapping particulates such as airborne pollutants on both bark and leaf hairs and are able to survive in very poor compacted ground. They can grow to a considerable size; although not the tallest, ‘Barnie’ was last measured as part of the Tree Register of the British Isles ([TROBI](#)) in 2017 and was then 35.2m tall and had a girth of 843cm.

Planes are the only Genus in the *Platanaceae* family, and the number of species recently increased from six to eleven, many from North America. Planes are not native to the UK, but Oriental plane is found across south-east Europe, Asia, and India. All but one species has simple palmately lobed (hand-shaped, fig. 2) leaves arranged alternately along the shoot (fig. 3) with an enlarged petiole (leaf stalk) base that encloses the associated bud on the shoot (fig.3). The flowers are wind-pollinated and borne in a ball-like structure, as are the seeds that follow, and these can remain on the tree for some time after leaves have fallen (fig. 3). Mature trees have distinctive bark which sheds in patches to give an attractive mottled appearance (fig. 3) though on old trees the trunk develops a more evenly coloured appearance.



Figure 1: Ravenscourt Park pot-bellied plane. Crown copyright, Forest Research

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations



Figure 2: Left to right - leaves of Oriental plane, London plane and Western plane (Westonbirt Arboretum)




Figure 3: Left to right - alternate leaf arrangement and swollen petiole bases, typical bark (© Crown copyright. Forestry Commission) and pom-pom fruits of London plane (Westonbirt Arboretum).

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations

Priority disease – Canker stain of plane (*Ceratocystis platani*)

Formerly known as Plane wilt, Canker stain of plane is a fungal disease of Oriental, Western, and London planes. It affects the water transport systems of the tree causing wilting of leaves and shoots and is usually fatal within 5-7 years. The fungus is native to North America and was first found in Europe in Marseille, France, in 1972. It is now present as far north as the Ile de France region of France, Italy, Switzerland, Albania, Greece, Turkey and Armenia. It has been successfully eradicated following outbreaks in Spain and Corsica.

The most visible symptom of infection is the sudden wilting of leaves, often initially on a single branch but soon spreading (fig. 4). As the infection progresses and spreads the crown of an infected tree becomes sparse and thin with discoloured leaves. Cankers can also form though they may not be readily visible, particularly on trees with thick bark. They typically present as sunken patches with orange and purple staining around their edges and beneath the bark as dark brown or violet spots of dead tissue (fig. 4), sometimes accompanied by more widespread blue-black staining. Dousing the bark with water can help to see their extent/features once detected.



Figure 4: Canker stain of plane symptoms (left to right) – isolated wilting/dead foliage (P. Tsopelas & N. Souiliorti, FIRA, Greece), sunken patch (canker) with typical peripheral orange-purple colouring (© Crown copyright. Forest Research), and exposed dark staining (lesion) beneath the bark (© Crown copyright. Forest Research).

Lookalikes

Western plane is particularly susceptible to **Plane Anthracnose** (*Apiognomonina veneta*), Oriental plane is moderately tolerant and susceptibility in London plane varies depending upon clone. The most visible symptom is the leaf-blight which presents as brown staining along the veins of leaves. Closer inspection will usually reveal cankers on the twigs or larger branches, shoot blight follows if these girdle stems (fig. 5). Frost and salt damage

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations

can give similar symptoms but in those cases twig cankers will be absent. Unlike Canker stain this fungus is not usually fatal.



Figure 5: Plane anthracnose symptoms (left to right) – leaf blight with brown staining along veins, twig blight with wilted/dead leaves, and deformity (canker) on a young twig (all images © Crown copyright. Forestry Commission).

Like other mildews **Plane powdery mildew** (*Erysiphe platani*) presents as white powdery patches on the leaves and shoots (fig. 6). Affected shoots and leaves go on to become distorted and turn yellow, eventually falling prematurely.

London plane trees are prone to infection by **Elbow-patch crust** (*Fomitoporia mediterranea*), a decay fungus which causes difficult to see cankers on the bark of affected trees (fig. 6). The associated dieback can look very similar to that caused by Plane wilt.



Figure 6: (L-R) Plane powdery mildew symptoms, large roughened deformity (canker) on a plane tree trunk caused by *Fomitoporia*, and a *Fomitoporia* fruit body (all images © Crown copyright. Forest Research).

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations

Priority pest – Plane lace bug (*Corythucha ciliata*)

Plane lace bug is native to North America and was accidentally introduced to mainland Europe. A small population was found at a plant nursery in Bedfordshire in 2006 but was successfully eradicated. There were no further reports until August 2024 when they were found on a number of urban trees in London after a passenger on a London bus found one on their rucksack. The lifecycle starts with eggs which are laid along the veins on the underside of plane leaves. The nymphs that hatch from the eggs are shiny black and oval shaped with a scattering of black spines and tend to stay clustered on the same leaf. After 5 moults they reach their adult form, around 4mm long, with white lace-like wing covers with some brown colouring at the base (fig. 7). Several generations can be produced each year and both adults and nymphs are often found together on the underside of leaves where they feed on the sap, usually with abundant dark spots of fresh and dried excreta (frass) (fig. 7).



Figure 7: Left to right - foliage spotting and discoloration caused by lace bugs (© Crown copyright. Forest Research), adult lace bug (Fera Science Ltd. Davis Cossley), and adults and nymphs with spots of frass on the underside of a plane leaf (Fera Science Ltd.).

Symptoms

Besides the presence of adult lace bugs, their nymphs, and their dark spots of frass, the main sign of lace bug presence is the appearance of small pale green or yellowish spots on the upper surface of leaves, usually clustered along the leaf veins. With increasing numbers of lace bugs the spots gradually coalesce resulting in whole leaves taking on a chlorotic pale-yellow colour with areas of bronzing. Very heavy infestations will lead to premature leaf drop and can, after several years, kill younger trees.

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations

Lookalikes

Leafhoppers are common sap-feeding insects that can produce symptoms very similar to the initial spotting/mottling caused by lace bugs. They are also found on the underside of leaves, but the insects are a different shape, lack the lacy appearance of adult lace bugs and readily take flight when disturbed (fig. 8).



Figure 8: Typical leaf hopper related damage on Oriental plane, and leaf hoppers on the underside of a sycamore leaf (Fera Science Ltd.)

Reporting

Canker stain of plane and Plane lace bug are both Observatree priorities so please report possible sightings via [TreeAlert](#).

Check the Observatree website for more information and resources on [Plane lace bug](#) and [Canker stain of plane](#). For more information about these and other pathogens affecting plane trees see [Pathology Advisory Note 7](#) from Forest Research.

Matt Parratt, Forest Research, August 2026

Observatree is a citizen science project led by Forest Research, in collaboration with key organisations